

TABLE 4.1 Anthropometric Data

| Segment | Definition | Segment Weight/Total | Center of Mass/ Segment Length | | Radius of Gyration/ Segment Length | | | Density |
|-----------------------------|--|----------------------|-----------------------------------|----------|---------------------------------------|--------|----------|---------|
| | | | Body Weight | Proximal | Distal | C of G | Proximal | |
| Hand | Wrist axis/knuckle II middle finger | 0.006 M | 0.506 | 0.494 P | 0.297 | 0.587 | 0.577 M | 1.16 |
| Forearm | Elbow axis/ulnar styloid | 0.016 M | 0.430 | 0.570 P | 0.303 | 0.526 | 0.647 M | 1.13 |
| Upper arm | Glenohumeral axis/elbow axis | 0.028 M | 0.436 | 0.564 P | 0.322 | 0.542 | 0.645 M | 1.07 |
| Forearm and hand | Elbow axis/ulnar styloid | 0.022 M | 0.682 | 0.318 P | 0.468 | 0.827 | 0.565 P | 1.14 |
| Total arm | Glenohumeral joint/ulnar styloid | 0.050 M | 0.530 | 0.470 P | 0.368 | 0.645 | 0.596 P | 1.11 |
| Foot | Lateral malleolus/head metatarsal II | 0.0145 M | 0.50 | 0.50 P | 0.475 | 0.690 | 0.690 P | 1.10 |
| Leg | Femoral condyles/medial malleolus | 0.0465 M | 0.433 | 0.567 P | 0.302 | 0.528 | 0.643 M | 1.09 |
| Thigh | Greater trochanter/femoral condyles | 0.100 M | 0.433 | 0.567 P | 0.323 | 0.540 | 0.653 M | 1.05 |
| Foot and leg | Femoral condyles/medial malleolus | 0.061 M | 0.606 | 0.394 P | 0.416 | 0.735 | 0.572 P | 1.09 |
| Total leg | Greater trochanter/medial malleolus | 0.161 M | 0.447 | 0.553 P | 0.326 | 0.560 | 0.650 P | 1.06 |
| Head and neck | C7-T1 and 1st rib/ear canal | 0.081 M | 1.000 | — PC | 0.495 | 0.116 | — PC | 1.11 |
| Shoulder mass | Sternoclavicular joint/glenohumeral axis | — | 0.712 | 0.288 | — | — | — | 1.04 |
| Thorax | C7-T1/T12-L1 and diaphragm* | 0.216 PC | 0.82 | 0.18 | — | — | — | 0.92 |
| Abdomen | T12-L1/L4-L5* | 0.139 LC | 0.44 | 0.56 | — | — | — | — |
| Pelvis | L4-L5/greater trochanter* | 0.142 LC | 0.105 | 0.895 | — | — | — | — |
| Thorax and abdomen | C7-T1/L4-L5* | 0.355 LC | 0.63 | 0.37 | — | — | — | — |
| Abdomen and pelvis | T12-L1/greater trochanter* | 0.281 PC | 0.27 | 0.73 | — | — | — | 1.01 |
| Trunk | Greater trochanter/glenohumeral joint* | 0.497 M | 0.50 | 0.50 | — | — | — | 1.03 |
| Trunk head neck | Greater trochanter/glenohumeral joint* | 0.578 MC | 0.66 | 0.34 P | 0.503 | 0.830 | 0.607 M | — |
| Head, arms, and trunk (HAT) | Greater trochanter/glenohumeral joint* | 0.678 MC | 0.626 | 0.374 PC | 0.496 | 0.798 | 0.621 PC | — |
| HAT | Greater trochanter/mid rib | 0.678 | 1.142 | — | 0.903 | 1.456 | — | — |

*NOTE: These segments are presented relative to the length between the greater trochanter and the glenohumeral joint.

Source Codes: M, Dempster via Miller and Nelson; *Biomechanics of Sport*, Lea and Febiger, Philadelphia, 1973. P, Dempster via Plagenhoef; *Patterns of Human Motion*, Prentice-Hall, Inc. Englewood Cliffs, NJ, 1971. L, Dempster via Plagenhoef from living subjects; *Patterns of Human Motion*, Prentice-Hall, Inc., Englewood Cliffs, NJ, 1971. C, Calculated.